

Net Weight lb (kg) on bag or bulk
Monensin Medicated Replacement Chicken Feed
Type C Medicated Feed

To aid in the prevention of coccidiosis caused by *Eimeria necatrix*, *E. tenella*, *E. acervulina*, *E. brunetti*, *E. mivati*, and *E. maxima*.

ACTIVE DRUG INGREDIENT

Monensin, USP90-110 g/ton*

GUARANTEED ANALYSIS

Crude Protein, not less than.....%
Lysine, not less than.....%
Methionine, not less than.....%
Crude Fat, not less than.....%
Crude Fiber, not more than.....%
Calcium, not less than.....%
Calcium, not more than.....%
Phosphorus, not less than.....%
Salt¹, not less than.....%
Salt¹, not more than.....%
Sodium², not less than.....%
Sodium², not more than.....%

¹If added.

²Shall be guaranteed only when total sodium exceeds that furnished by the maximum salt guarantee.

INGREDIENTS

Each ingredient must be named in accordance with the names and definitions adopted by the Association of American Feed Control Officials.

FEEDING DIRECTIONS

Feed 90-110 g/ton of complete feed to replacement chickens intended for use as cage layers. Feed continuously as the sole ration.

CAUTION

Do not feed to laying chickens. Do not feed to chickens over 16 weeks of age. Do not allow horses, other equines, mature turkeys, or guinea fowl access to feed containing monensin. Ingestion of monensin by horses and guinea fowl has been fatal.

MANUFACTURED BY

BLUE BIRD FEED MILL
Any town, USA 12345

*Final printed label on formulated Type C medicated feed must bear a single drug concentration.