

Net Weight lb (kg) on bag or bulk
Monensin Medicated Turkey Feed
Type C Medicated Feed

For the prevention of coccidiosis caused by *Eimeria adenoeides*, *E. meleagrimitis*, and *E. gallopavonis*.

ACTIVE DRUG INGREDIENT

Monensin, USP54-90 g/ton*

GUARANTEED ANALYSIS

Crude Protein, not less than.....	_____	%
Lysine, not less than.....	_____	%
Methionine, not less than.....	_____	%
Crude Fat, not less than.....	_____	%
Crude Fiber, not more than.....	_____	%
Calcium, not less than.....	_____	%
Calcium, not more than.....	_____	%
Phosphorus, not less than.....	_____	%
Salt ¹ , not less than.....	_____	%
Salt ¹ , not more than.....	_____	%
Sodium ² , not less than.....	_____	%
Sodium ² , not more than.....	_____	%

¹If added.

²Shall be guaranteed only when total sodium exceeds that furnished by the maximum salt guarantee.

INGREDIENTS

Each ingredient must be named in accordance with the names and definitions adopted by the Association of American Feed Control Officials.

FEEDING DIRECTIONS

Feed 54-90 g/ton of complete feed. Feed continuously as the sole ration.

CAUTION

For growing turkeys only. Do not allow horses, other equines, mature turkeys, or guinea fowl access to feed containing monensin. Ingestion of monensin by horses and guinea fowl has been fatal. Some strains of turkey coccidia may be monensin tolerant or resistant. Monensin may interfere with development of immunity to turkey coccidiosis.

MANUFACTURED BY

BLUE BIRD FEED MILL
Any town, USA 12345

*Final printed label on formulated Type C medicated feed must bear a single drug concentration.