

**Net Weight lb (kg) on bag or bulk**  
**Ractopamine Medicated Tom Turkey Feed**  
**Type B Medicated Feed**

**TOMS:** For increased rate of weight gain and improved feed efficiency in finishing tom turkeys when fed for the last 14 days prior to slaughter.

**ACTIVE DRUG INGREDIENT**

Ractopamine Hydrochloride.....11.9 to 4920 g/ton\*

**GUARANTEED ANALYSIS**

Crude Protein, not less than.....	_____	%
Lysine, not less than.....	_____	%
Methionine, not less than.....	_____	%
Crude Fat, not less than.....	_____	%
Crude Fiber, not more than.....	_____	%
Calcium, not less than.....	_____	%
Calcium, not more than.....	_____	%
Phosphorus, not less than.....	_____	%
Salt <sup>1</sup> , not less than.....	_____	%
Salt <sup>1</sup> , not more than.....	_____	%
Sodium <sup>2</sup> , not less than.....	_____	%
Sodium <sup>2</sup> , not more than.....	_____	%

<sup>1</sup> If added.

<sup>2</sup> Shall be guaranteed only when total sodium exceeds that furnished by the maximum salt guarantee.

**INGREDIENTS**

Each ingredient must be named in accordance with the names and definitions adopted by the Association of American Feed Control Officials.

**MIXING DIRECTIONS**

Thoroughly mix ractopamine Type B Medicated Feed into one ton of complete turkey feed, according to the table below to obtain the proper concentration in the Type C Medicated Feed. Prepare an intermediate pre-blend of the medicated feed prior to mixing in a complete feed. Thoroughly mix the required amount in a convenient quantity of feed ingredients, then add to the remaining feed ingredients to make a ton of complete feed. [Use only the portion of the table below on your Type B Medicated Feed product label that is applicable to the concentration of ractopamine in the Type B Medicated Feed which you manufacture. The following table gives examples of how some Type B Medicated Feed concentrations can be prepared.]

\*Final printed label on formulated Type C medicated feed must bear a single drug concentration. No increased benefit has been shown when ractopamine concentrations in the diet are greater than 4.6 g/ton (5 ppm)

Ractopamine Concentration in Type B Feed (g/ton)	Desired Ractopamine Concentration in Type C Medicated Feed (g/ton)**		
	4.6 g/ton	8.2 g/ton	11.8 g/ton
	lb of Type B per ton of Ractopamine Medicated Complete Feed		
20	460	820	1,180
200	46	82	118
1,000	9.2	16.4	23.6

\*\* No increased benefit has been show when ractopamine concentrations in the diet are greater than 4.6 g/ton (5 ppm).

### CAUTION

Feeding ractopamine to tom turkeys during periods of excessive heat can result in increased mortality. Not for animals intended for breeding.

### WARNING

The active ingredient in Topmax, ractopamine hydrochloride, is a beta-adrenergic agonist. Individuals with cardiovascular disease should exercise special caution to avoid exposure. Not for use in humans. Keep out of the reach of children. The Topmax 9 formulation (Type A Medicated Article) poses a low dust potential under usual conditions of handling and mixing. When mixing and handling Topmax, use protective clothing, impervious gloves, protective eye wear, and a NIOSH-approved dust mask. Operators should wash thoroughly with soap and water after handling. If accidental eye contact occurs, immediately rinse eyes thoroughly with water. If irritation persists, seek medical attention. The material safety data sheet contains more detailed occupational safety information. To report adverse effects, access medical information, or obtain additional product information, call 1-800-428-4441.

**MANUFACTURED BY**  
**BLUE BIRD FEED MILL**  
 Any town, USA 12345